COPD Pharmacotherapy Review

Based on the 2025 GOLD Report (Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease)

pyrls

More clinical pearls at pyrls.com

Abbreviations

eos Blood eosinophil count (cells/mcL)
ICS Inhaled corticosteroid
LABA Long-acting beta-agonist
LAMA Long-acting muscarinic antagonist
SABA Short-acting beta-agonist
SAMA Short-acting muscarinic antagonist

COPD Treatment Goals

Stable COPD	COPD Exacerbations
• **Improve** symptoms, exercise tolerance, health	• **Minimize** the effects of the exacerbation
• **Reduce risks of** disease progression, exacerbations, death	• **Prevent** future exacerbations

Initiation of Pharmacotherapy in Stable COPD

Select initial pharmacotherapy based on current symptoms and exacerbation history

Note
- If combination therapy is chosen, consider **single-inhaler options** for patient convenience
- Short-acting bronchodilators should be prescribed to all patients for immediate symptom relief

Exacerbations

≥2 moderate exacerbations OR ≥1 resulting in hospital admission

E

LABA + LAMA

- If eos ≥300, consider LABA + LAMA + **ICS**

≤1 moderate exacerbations not resulting in hospital admission

A

Bronchodilator
(LABA, LAMA, SABA, or SAMA)

- Either long- or short-acting, but **long-acting is generally preferred** unless very occasional dyspnea

CAT <10, mMRC 0-1

B

LABA + LAMA

- If LABA + LAMA combo is not appropriate, **choose either** based on the patient's perceived symptom relief

CAT ≥10, mMRC ≥2

Symptoms

CAT: COPD Assessment Test
mMRC: Modified Medical Research Council dyspnea scale

Adapted from Figure 3.7 of the 2025 GOLD Report

Pharmacotherapy Key Points

General Notes on Inhaled Medications

- **Individualize** inhaler choice (e.g., cost, ability to use, efficacy)
- Assess inhaler technique and adherence prior to therapy modification

Bronchodilators - First-Line for COPD

- **Long-acting** agents (LABA, LAMA) are **preferred** over short-acting (SABA, SAMA), except in those with only **occasional dyspnea** and when **immediate relief** is needed in those on long-acting maintenance therapy
- **Combination** of LAMA+LABA is more effective at improving symptoms and lung function than monotherapy; patients not managed on a single long-acting bronchodilator should be **escalated** to dual therapy
- LAMAs provide greater exacerbation risk reduction than LABAs
- Combination of SABA+SAMA is more effective than either alone
- **Inhaled** bronchodilators are recommended over oral bronchodilators

Inhaled Corticosteroids (ICS) - Main Anti-Inflammatory Therapy

- ICS **monotherapy** has **not** shown long-term benefit in COPD treatment
- Can be **added** to LABA+LAMA in patients with exacerbation history (especially if recurrent or has led to hospitalization) or **eos ≥100**
  - ICS **should** be included if features of **asthma** are present
- ICS+LABA+LAMA has proven mortality benefit versus LABA+LAMA in those with symptomatic COPD and history of exacerbations
- ICS+LABA **not** encouraged for COPD; if ICS is indicated, use **triple regimen**
- Main safety concern is **pneumonia**; not recommended in patients with history of **recurrent pneumonia** or **mycobacterial infection**

Others - PDE Inhibitors, Dupilumab, Antibiotics

- **Roflumilast** (PDE-4 inhibitor) reduces **exacerbations** and improves **lung function** in those with **severe to very severe airflow limitation**, **chronic bronchitis**, and **exacerbations**
  - Consider adding to LABA+LAMA (+/- ICS) for these patients
- **Ensifentrine** (PDE-3 and -4 inhibitor) improves lung function and dyspnea
  - Consider adding to LABA+LAMA in patients with **dyspnea**
- **Dupilumab** (IL-4 and IL-13 inhibitor) reduces exacerbations and improves lung function in patients with **chronic bronchitis**, history of **exacerbations** despite LABA+LAMA+ICS therapy, and **eos ≥300**
  - Consider adding to triple therapy for these patients if exacerbations occur
- In those with exacerbations despite appropriate therapy, **macrolides** (e.g., azithromycin) may be considered (especially in **former smokers**)

Follow-Up Pharmacotherapy Management in Stable COPD

COPD management is an individualized, continuous cycle of assessment and treatment adjustment

Is COPD well managed?

Review:
- Symptoms (e.g., dyspnea)
- Exacerbations

Assess:
- Inhaler technique & adherence
- Non-pharmacological interventions

Adjust:
- Consider escalation/de-escalation
- Switch device or medications

No →

If Yes: Continue current therapy

Adapted from Figures 3.8 and 3.9 of the 2025 GOLD Report

Inhalers Chart + Administration Guides

Primary Issue?*

Dyspnea

LAMA or LABA

↓

LAMA + LABA**

↓

- Consider switching inhaler or medication
- Optimize non-pharmacological interventions
- Assess and address other causes of symptoms
- Consider adding **ensifentrine**

Exacerbations

LAMA or LABA

↓

LAMA + LABA**

← eos <300 / eos ≥300 ↓

eos ≥100 / eos <100 ↓

LAMA + LABA + ICS**
(Consider **de-escalating** to LAMA + LABA if ICS is not tolerated or not effective)

→ If eos ≥300 ↓ Consider adding **dupilumab** if symptoms of **chronic bronchitis** are present

Consider adding:
- **Roflumilast** if FEV1 <50% + chronic bronchitis
- **Azithromycin**, preferentially in former smokers

Although LABA+ICS is not preferred in treatment of COPD without features of asthma, if the patient has already been on LABA+ICS and is well managed, the current therapy **may** be continued (or consider switching to LABA+LAMA if no relevant exacerbation history).
- If **exacerbations** occur, escalate to LABA+LAMA+ICS (if eos ≥100) **or** switch to LABA+LAMA (if eos <100)
- If major symptoms are present, decide based on previous response to ICS:
  - If no relevant history of exacerbation, consider switching to LABA+LAMA
  - If history of positive response to ICS in previous exacerbations, consider escalating to LABA+LAMA+ICS

*If **both** dyspnea and exacerbation must be addressed, **use the exacerbation pathway**
**For patients on LAMA+LABA or LAMA+LABA+ICS, single-inhaler options should be considered for convenience

Pharmacotherapy Management of Acute Exacerbations*

*Non-life-threatening

Initial Treatment: SABA (with or without SAMA) →

- Initiate maintenance with **long-acting bronchodilators** as soon as stable
  - Consider adding **ICS** to LAMA+LABA if frequent exacerbations with ↑ eos
- If severe exacerbation, consider **systemic corticosteroids** (duration: generally **≤5 days**)
- If indicated (e.g., signs of bacterial infection), give **antibiotics** (duration: generally **5-7 days**)

Reference: Global Initiative for Chronic Obstructive Lung Disease. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease (2025 Report)

® 2024 Cosmas Health, Inc. and/or its affiliates. All rights reserved.

Last Revised 11/2024